

AD-A055 139

STANFORD UNIV CALIF INST FOR PLASMA RESEARCH  
RESISTIVE DECAY OF PHOTOSPHERIC CURRENT SHEETS.(U)  
MAY 78 D D BARBOSA

F/G 4/1

UNCLASSIFIED

SU-IPR-744

N00014-75-C-0673

NL

| OF |  
AD  
A055139



END  
DATE  
FILMED  
7-78  
DDC

FOR FURTHER TRAN *Dr. A. M. 1978*

AD A 055139

# RESISTIVE DECAY OF PHOTOSPHERIC CURRENT SHEETS

by  
D.D. Barbosa


12  
SC

National Aeronautics and Space Administration  
NGL 05-020-272

and

Office of Naval Research  
Contract N00014-75-C-0673

SUIPR Report No. 744



May 1978

*gfb* E

**DISTRIBUTION STATEMENT A**

Approved for public release;  
Distribution Unlimited



**INSTITUTE FOR PLASMA RESEARCH  
STANFORD UNIVERSITY, STANFORD, CALIFORNIA**

78 06 14 045

AD No. \_\_\_\_\_  
DDC FILE COPY

RESISTIVE DECAY OF PHOTOSPHERIC CURRENT SHEETS



by

D. D. Barbosa

Institute for Plasma Research  
Stanford University  
Stanford, California 94305

ABSTRACT

We consider the magnetostatics of thin, axi-symmetric current sheets in the plane of the photosphere. The resistive decay time constant for energy and flux of the system is computed and found to depend linearly on the thickness of the current sheet.





## I. INTRODUCTION

The possibility that photospheric neutral winds can blow across solar magnetic fields and thereby drive photospheric and coronal electric currents has been considered by Sen and White (1972) and Heyvaerts (1974) [see also Obayashi (1975)]. The primary motivation of these authors for developing a dynamo theory was to model a current system which, after the onset of an instability, could be rapidly dissipated and provide sufficient energy for a flare. More recently, Barbosa (1978) has considered some of the more general aspects of photospheric dynamo action.

One important conclusion shared by all of these authors is that when a photospheric current system is confined to a thin sheet the thickness of the sheet plays a central role in almost every aspect. In particular, the resistive time constant of the system should depend on the thickness. The question is, just how?

In Section II we consider the magnetostatics of axi-symmetric, azimuthal current sheets and describe one simple case for illustrative purposes. In Section III we compute the resistive time constants (energy and flux) for these systems.

ACCESSION for		
NTIS	White Section	<input checked="" type="checkbox"/>
DDC	Buff Section	<input type="checkbox"/>
UNANNOUNCED		<input type="checkbox"/>
JUSTIFICATION.....		
BY.....		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL. and/or SPECIAL	
A		

## II. MAGNETOSTATICS

We consider current restricted to flow in a thin sheet at  $z=0$  [ $(\rho, \phi, z)$  coordinates] where the thickness and perpendicular scale length satisfy  $\ell \ll r$ . The current density is modelled as

$$\vec{J}(\vec{x}) = K(\rho) \frac{1}{\ell} \left[ H(z + \ell/2) - H(z - \ell/2) \right] \hat{\phi} \approx K(\rho) \delta(z) \hat{\phi}, \quad (1)$$

where  $H(z)$  is the Heaviside step function and  $K(\rho)$  is the surface current density in the sheet with scale length  $r$ . The magnetic vector potential for this system is given as [Jackson, 1962]

$$A_{\phi}(\vec{x}) = \frac{2\pi}{c} \int_0^{\infty} \rho' d\rho' K(\rho') \int_0^{\infty} dk J_1(k\rho) J_1(k\rho') e^{-k|z|} \quad (2)$$

and the field lines may be obtained from the Euler potential

$$\alpha = \rho A_{\phi} = \text{constant}. \quad (3)$$

The magnetic field in cylindrical coordinates is

$$B_z = \frac{1}{\rho} \frac{\partial \alpha}{\partial \rho}, \quad B_{\rho} = -\frac{1}{\rho} \frac{\partial \alpha}{\partial z}. \quad (4)$$

We shall make use of the Fourier-Bessel transformation throughout this paper

$$\left. \begin{aligned} \tilde{K}(k) &= \int_0^{\infty} \rho d\rho K(\rho) J_1(k\rho) \\ K(\rho) &= \int_0^{\infty} k dk \tilde{K}(k) J_1(k\rho) \end{aligned} \right\} \quad (5)$$

Thus, the vector potential is simply

$$A_{\phi} = \frac{2\pi}{c} \int_0^{\infty} dk \tilde{K}(k) J_1(k\rho) e^{-k|z|} , \quad (6)$$

and the total current flowing is

$$I = \int_0^{\infty} d\rho K(\rho) = \int_0^{\infty} dk \tilde{K}(k) . \quad (7)$$

Relatively simple expressions for the magnetic field may be obtained from the class of current densities given by

$$K(\rho) = K_0 \frac{r^{2n} \rho}{(\rho^2 + r^2)^{n+1/2}} . \quad (8)$$

Gleeson and Axford (1976) have considered this class of currents in connection with current sheets in the Jovian magnetosphere. We do not propose that they are accurate representations of a photospheric current system. They serve only as convenient models which possess easily describable features and for which our calculations may be performed with significant tractability.

The case of  $n=1$  is a monopolar field with straight field lines which behaves as  $B \sim \rho^{-2}$ . A more realistic case is given by  $n=2$  for which the field lines are derived from

$$\alpha = \frac{2\pi K_0 r^2}{3c} \frac{r\rho^2}{[\rho^2 + (|z|+r)^2]^{3/2}} = \frac{2\pi K_0 r^2}{3c} \bar{\alpha} . \quad (9)$$

This configuration has an O-type neutral point at  $\rho = \sqrt{2} r$  and the current links a maximum (normalized) flux there of  $\bar{\alpha}_{MAX} = 2/\sqrt{27}$ .



Representative field lines are shown in Figure 1. The actual magnetic flux distribution is given by

$$\Phi = 2\pi\alpha . \quad (10)$$

### III. RESISTIVE TIME CONSTANTS

To compute the resistive decay time, we assume that a given current system (1) has been set up and at  $t=0$  all electromotive forces are shut off and the system decays through ohmic losses. Integration over all space yields

$$E = \frac{1}{8\pi} \int d\vec{x} B^2 = \frac{1}{2c} \int d\vec{x} \vec{J} \cdot \vec{A} , \quad (11)$$

and making use of (5) we find that the total energy of the system is given by

$$E = \frac{2\pi^2}{c^2} \int_0^\infty dk \tilde{K}^2(k) . \quad (12)$$

The self-inductance of the system is defined by  $E = \frac{1}{2} LI^2$  and therefore

$$L = \frac{4\pi^2}{c^2} \frac{1}{I^2} \int_0^\infty dk \tilde{K}^2(k) . \quad (13)$$

Ohmic losses occur throughout the volume of the current sheet and Barbosa (1978) has shown that the power dissipated per unit area of the sheet is given as

$$\mathcal{P} = \frac{K^2(\rho)}{\sigma_3 \ell} , \quad (14)$$

where  $\sigma_3$  is the Cowling conductivity. Integration over the entire area of the sheet gives the total power dissipated and upon equating this to  $I^2 R$  we find that the effective resistance of the system is given by



$$R = \frac{2\pi}{\sigma_3} \frac{1}{l^2} \int_0^\infty k \, dk \, \tilde{K}^2(k) . \quad (15)$$

Elementary circuit theory gives time constants for decay of the energy and current (or flux) as

$$\tau_E = L/2R , \quad \tau_\Phi = L/R . \quad (16)$$

Thus, the time constant for decay of magnetic flux is (esu)

$$\tau_{\text{DIFF}} = \tau_\Phi = \frac{2\pi\sigma_3 l r}{c^2} g , \quad (17)$$

where  $g$  is a geometrical factor of order one,

$$g = \frac{1}{r} \frac{\int_0^\infty dk \, \tilde{K}^2(k)}{\int_0^\infty k \, dk \, \tilde{K}^2(k)} \quad (18)$$

The evaluation of  $g$  for the model of (8) yields  $g = 2, 2/3$  for  $n=1,2$  respectively.

Our basic result is that the decay time depends only linearly on the thickness  $l$  for the simple reason that magnetic energy is stored in a volume  $\sim r^3$  but ohmic losses occur in a volume  $\sim l r^2$ .

#### ACKNOWLEDGMENT

This work was supported by NASA Grant NGL 05-020-272 and by ONR Contract N00014-C-0673.

#### REFERENCES

- Barbosa, D. D. 1978, "Dynamo Action in a Thin Slab", submitted to  
Astrophys. J.
- Gleeson, L. J. and Axford, W. I. 1976, Jour. Geophys. Res., 81, 3403.
- Heyvaerts, J. 1974 Solar Phys., 38, 419.
- Jackson, J. D. 1962, Classical Electrodynamics, (New York: Wiley), p. 166.
- Obayashi, T. 1975, Solar Phys., 40, 217.
- Sen, H. K. and White, M. L. 1972, Solar Phys., 23, 146.

# RESISTIVE DECAY OF PHOTOSPHERIC CURRENT SHEETS

by


D. D. Barbosa

Institute for Plasma Research  
Stanford University  
Stanford, California 94305

## CAPTION FOR FIGURE

Figure 1. Field lines of a current sheet in the  $z=0$  plane for  $n=2$  in Equation (8). The normalized flux surfaces  $\bar{\alpha}$  satisfy Equation (9) and are displayed for  $\bar{\alpha} = 0.1$  to 0.35 in steps of 0.05.  $\bar{\alpha}_{\text{MAX}} \approx 0.38$  occurs at  $\rho = \sqrt{2} r$ .





REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER SUHPR- <u>                    </u> 744 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) RESISTIVE DECAY OF PHOTOSPHERIC CURRENT SHEETS,	5. TYPE OF REPORT & PERIOD COVERED Scientific, Technical	
7. AUTHOR(s) D.D. Barbosa	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Institute for Plasma Research Stanford University Stanford, California 94305	8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0673 VNG-L-05-020-272	
11. CONTROLLING OFFICE NAME AND ADDRESS Phil Surra, Office of Naval Research Durand 165, Stanford University	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Scientific rept.,	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE May 78	
	13. NUMBER OF PAGES 10	
	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)  This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Solar magnetism, Photospheric currents		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  We consider the magnetostatics of thin, axi-symmetric current sheets in the plane of the photosphere. The resistive decay time constant for energy and flux of the system is computed and found to depend linearly on the thickness of the current sheet.		